

## ORIGINAL ARTICLE

# Combined Immunodeficiency Associated with *DOCK8* Mutations

Qian Zhang, M.D., Jeremiah C. Davis, M.P.H., Ian T. Lamborn, B.S., Alexandra F. Freeman, M.D., Huie Jing, Ph.D., Amanda J. Favreau, B.S., Helen F. Matthews, B.S.N., Joie Davis, M.S.N., Maria L. Turner, M.D., Gulbu Uzel, M.D., Steven M. Holland, M.D., and Helen C. Su, M.D., Ph.D.

## ABSTRACT

**BACKGROUND**

Recurrent sinopulmonary and cutaneous viral infections with elevated serum levels of IgE are features of some variants of combined immunodeficiency. The genetic causes of these variants are unknown.

**METHODS**

We collected longitudinal clinical data on 11 patients from eight families who had recurrent sinopulmonary and cutaneous viral infections. We performed comparative genomic hybridization arrays and targeted gene sequencing. Variants with predicted loss-of-expression mutations were confirmed by means of a quantitative reverse-transcriptase–polymerase-chain-reaction assay and immunoblotting. We evaluated the number and function of lymphocytes with the use of in vitro assays and flow cytometry.

**RESULTS**

Patients had recurrent otitis media, sinusitis, and pneumonias; recurrent *Staphylococcus aureus* skin infections with otitis externa; recurrent, severe herpes simplex virus or herpes zoster infections; extensive and persistent infections with molluscum contagiosum; and human papillomavirus infections. Most patients had severe atopy with anaphylaxis; several had squamous-cell carcinomas, and one had T-cell lymphoma–leukemia. Elevated serum IgE levels, hypereosinophilia, low numbers of T cells and B cells, low serum IgM levels, and variable IgG antibody responses were common. Expansion in vitro of activated CD8 T cells was impaired. Novel homozygous or compound heterozygous deletions and point mutations in the gene encoding the dedicator of cytokinesis 8 protein (*DOCK8*) led to the absence of *DOCK8* protein in lymphocytes.

**CONCLUSIONS**

Autosomal recessive *DOCK8* deficiency is associated with a novel variant of combined immunodeficiency.

From the Laboratory of Host Defenses (Q.Z., I.T.L., H.J., H.C.S.), the Laboratory of Clinical Infectious Diseases (A.F.F., J.D., G.U., S.M.H.), the Research Technologies Branch (A.J.F.), and the Laboratory of Immunology (H.F.M.), National Institute of Allergy and Infectious Diseases; the Howard Hughes Medical Institute (J.C.D.); and the Dermatology Branch, National Cancer Institute (M.L.T.) — all in Bethesda, MD. Address reprint requests to Dr. Su at the National Institute of Allergy and Infectious Diseases, National Institutes of Health, CRC 5W3940, MSC 1456, Bethesda, MD 20892, or at [hsu@niaid.nih.gov](mailto:hsu@niaid.nih.gov).

This article (10.1056/NEJMoa0905506) was published on September 23, 2009, and updated on October 20, 2009, at [NEJM.org](http://NEJM.org).

N Engl J Med 2009;361:2046-55.  
Copyright © 2009 Massachusetts Medical Society.

**I**NHERITED SUSCEPTIBILITY TO VIRAL INFECTIONS, especially by herpesviruses, occurs in patients who have developmental defects of T cells, such as severe combined immunodeficiency; disorders involving cell killing by cytotoxic lymphocytes, such as hemophagocytic lymphohistiocytosis and X-linked lymphoproliferative disease; and impaired type I interferon production.<sup>1,2</sup> Several other conditions confer susceptibility to human papillomavirus infections or molluscum contagiosum.<sup>3-6</sup>

The hallmarks of severe combined immunodeficiency are impaired T-cell function and a severe deficiency of T cells. In some forms of the disease, there are also low numbers of or dysfunctional B cells and natural killer cells. The molecular defects in severe combined immunodeficiency are diverse,<sup>1,7-13</sup> and forms of the disease that are caused by residual gene activity or partial T-cell immunodeficiencies (e.g., the Wiskott–Aldrich syndrome) can feature skin diseases with elevated levels of IgE and hypereosinophilia.<sup>14,15</sup>

We describe 11 patients who had a combined immunodeficiency with low numbers of T, B, and natural killer cells, extensive cutaneous viral infections, and susceptibility to cancer. These patients had loss-of-function mutations in an uncharacterized gene, dedicator of cytokinesis 8 (*DOCK8*), which is expressed in lymphocytes.

## METHODS

### CELLS

We isolated subgroups of T cells from peripheral-blood mononuclear cells by negative selection, using magnetic bead isolation and then stimulation with anti-CD3 plus anti-CD28 antibodies (1  $\mu$ g per milliliter) and cultured in interleukin-2 (for details, see the Supplementary Appendix, available with the full text of this article at NEJM.org). B cells were immortalized with the Epstein–Barr virus, and T cells with herpesvirus saimiri (HVS), according to standard protocols.

### COMPARATIVE GENOMIC HYBRIDIZATION ANALYSES

Comparative genomic hybridization analyses were performed with 244K arrays with the use of the Agilent platform, according to the manufacturer's instructions. Modifications of the process are described in the Supplementary Appendix.

### DNA SEQUENCING

Sequencing of genomic DNA was performed after polymerase-chain-reaction (PCR) amplification of exons with their flanking intronic or untranslated regions. (Primer sequences are listed in Table 1 in the Supplementary Appendix.) Point mutations were confirmed in a second PCR amplification reaction, and frameshift mutations were confirmed after cloning. Novel variants that were identified in the 11 index patients were sought in 6 patients with autosomal dominant hyperimmunoglobulin E syndrome (HIES), in 32 patients with other immunologic diseases, in 15 healthy blood donors of various ethnic backgrounds, and in 100 healthy white control subjects.

### IMMUNOBLOTTING

Cytosolic lysates were prepared according to standard methods, followed by separation on 3 to 8% TRIS-acetate gels (Invitrogen) and transfer onto nitrocellulose. Proteins were detected with the use of polyclonal rabbit anti-DOCK8 antibodies (Sigma-Aldrich or donated by Dr. Ruusala, Ludwig Cancer Institute, Uppsala, Sweden<sup>16</sup>) or mouse anti- $\beta$ -actin antibodies (Sigma-Aldrich) on the same blot. (For details, see the Methods section in the Supplementary Appendix.)


### FLOW CYTOMETRY

Standard flow-cytometric methods were used for staining of cell-surface markers and carboxyfluorescein succinimidyl ester (CFSE) dilution assays. Cells were harvested for CFSE analysis after 3 days of culture under conditions of stimulation or no stimulation with anti-CD3 plus anti-CD28 antibodies. (See the Supplementary Appendix for additional details.)

## RESULTS

### CLINICAL FINDINGS

The patients and their relatives were enrolled in research protocols that were approved by the institutional review board of the National Institute of Allergy and Infectious Diseases. All study subjects provided written informed consent. The clinical features of Patient 1 in Family 3, Patients 1 and 2 in Family 4, and Patients 1 and 2 in Family 8 have been described previously.<sup>5,17,18</sup> We initially studied three patients (the index patients from



**Figure 1. Characteristic Dermatologic Findings.**

Among the characteristic dermatologic findings in the patients in the study were atopic dermatitis with scattered superimposed molluscum contagiosum lesions on the inside of the arm (Panel A), chronic ulcerative anogenital herpes simplex virus infection (Panel B), periungual and acral warts (Panel C), and molluscum contagiosum on the back (Panel D).

Families 1, 2, and 3) from a group of patients with undefined combined immunodeficiencies. After finding that the three carried a mutation in the same gene (*DOCK8*), we recognized that their disorders shared certain clinical features with atypical forms of HIES. We therefore sought out additional patients and included in our study several patients who were described previously, for a total of 11 patients between the ages of 6 and 21 years from eight families<sup>5,18</sup> (Table 2 in the Supplementary Appendix). Both male and female patients of various ethnic backgrounds were affected. Two patients presented with rash at birth, and all had atopic dermatitis (Fig. 1A). Of the 11 patients, 9 had severe and extensive food or environmental allergies, including anaphylaxis, and 6 had asthma or reactive airway disease. Eosinophilic esophagitis or lung disease was observed in two patients.

All the patients had recurrent upper and lower respiratory tract infections. Most of the patients had otitis media that had required placement of a tympanostomy tube, two patients had had mastoiditis, and seven patients had recurrent sinusitis. Nine patients had recurrent pneumonias or bronchitis, and bronchiectasis had been diagnosed in two patients. Recovered pulmonary pathogens included *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Pneumocystis jiroveci*, respiratory adenovirus, and respiratory syncytial virus. One patient had recurrent croup.

All the patients had extensive, frequently co-existing, cutaneous viral infections. Superficial, often ulcerating, herpes simplex virus infection occurred in seven patients, manifested as recurrent or persistent orolabial or anogenital involvement, keratitis, or eczema herpeticum (Fig. 1B). Seven patients had persistent flat and verrucous warts (Fig. 1C); five had extensive, disfiguring molluscum contagiosum infections (Fig. 1D); two had recurrent herpes zoster; and one had severe primary varicella-zoster infection.

Eight patients had *Staphylococcus aureus* skin infections or abscesses, two had *S. aureus* osteomyelitis, five had mucosal or nail candidiasis, and six had recurrent otitis externa. One patient had had cryptococcal and *H. influenzae* meningitis. Other infections included several occurrences of salmonella enteritis, giardiasis, and pericarditis.

Except for Patient 1 in Family 8, the 11 patients we studied did not have the neurologic, vasculitic, or autoimmune findings reported in patients with autosomal recessive HIES.<sup>5</sup> Nonimmuno-



logic features that are typically seen in autosomal dominant HIES were uncommon.<sup>19</sup> Three patients had poor growth, and one had delayed puberty.

Vulvar, facial, and anal squamous-cell dysplasia and carcinomas had occurred in patients with long-standing herpes simplex virus infections, human papillomavirus infections, and molluscum contagiosum. Cancers had developed in three patients during late childhood or early adulthood. Two patients died from metastatic squamous-cell carcinoma. One patient with resected microcystic adenoma died from cutaneous T-cell lymphoma–leukemia.<sup>17</sup>

### IMMUNOLOGIC ASSESSMENT

Studies of peripheral-blood mononuclear cells revealed low absolute lymphocyte counts in 9 of the 11 patients, including low counts for total T cells (in 10 of 11 patients), CD4 T cells (in all 11 patients), and CD8 T cells (in 10 of 11 patients) (Fig. 2).<sup>20–22</sup> Ratios of T4 cells to T8 cells were within normal ranges. The numbers of regulatory T cells, as assessed by CD4+CD25<sup>hi</sup>FOXP3+ coexpression, in two patients were decreased because of overall lymphopenia but were proportionally normal (Fig. 1 in the Supplementary Appendix). The number of natural killer cells was decreased in 6 of 10 patients, and the number of B cells was decreased in 5 of 11 patients. Although one patient had a normal number of eosinophils, most patients had mild-to-moderate eosinophilia, with a mean ( $\pm$ SD) of  $2.021 \pm 1.292 \times 10^3$  cells per cubic millimeter (normal count,  $<0.600 \times 10^3$ ). Patient 1 in Family 8 had an eosinophil count as high as  $33 \times 10^3$  per cubic millimeter. The numbers of neutrophils and monocytes were normal in all patients (data not shown).

Despite a decreased number of B cells in some patients, six patients had hypergammaglobulinemia, and five had normal levels of serum IgG. Levels of serum IgA varied, but levels of IgM were consistently low in all patients, with a mean of  $35 \pm 13$  mg per deciliter (normal value,  $>49$ ). Except for Patient 2 in Family 5, who had high-normal IgE levels (up to 818 IU per milliliter), the patients had very high IgE levels (peak range, 5630 to 43,600 IU per milliliter) (Fig. 2).


Levels of antibodies against a panel of bacterial and viral antigens were variable (Table 3 in the Supplementary Appendix). All patients who were evaluated had protective levels of antibodies

to rubella (8 of 8 patients) and varicella–zoster virus (4 of 4 patients). Some patients had a response to vaccines against *H. influenzae* type B (4 of 9 patients), diphtheria toxoid (5 of 10 patients), and tetanus toxoid (3 of 11 patients). Patients 1 and 2 in Family 4 had impaired T-cell–dependent primary antibody responses after immunization with the neoantigen bacteriophage  $\phi$ X174, with peak rates of bacteriophage inactivation in serum of 0.8 and 0.4 per minute, respectively, at 7 days (normal range at 14 days, 7 to 500). These two siblings also did not have an IgG antibody response after a second challenge with the bacteriophage antigen, with peak rates of inactivation of 0.7 and 0.3 per minute, respectively, on the day of the second challenge with no IgG. (The normal rate of inactivation in serum peaks at 7 days after a second challenge, with a range of 700 to 2000 per minute and with a mean IgG representing  $56 \pm 24\%$  of the neutralizing immunoglobulins.) Responses to pneumococcal polysaccharide antigens were variable (Table 3 in the Supplementary Appendix). Six of the patients were treated with long-term replacement immune globulin, which seemed to improve the incidence and severity of sinopulmonary infections in three patients without influencing the course of viral infections in any patient.

### GENETIC ANALYSES

Figure 3A shows the pedigrees of the eight families. Families 1, 2, and 8 were consanguineous. Comparative genomic hybridization arrays of genomic DNA from the index patients in Families 1 and 2 revealed homozygous deletions in the *DOCK8* gene (Fig. 3B, and Fig. 2A in the Supplementary Appendix). Deletion A in Family 1 spanned exons 10 through 23, and deletion B in Family 2 spanned exons 5 through 24 (Fig. 3B, and Fig. 2A and 2C in the Supplementary Appendix). Deletions A and B were confirmed by the failure of PCR to amplify deleted exons; furthermore, the juxtaposition of normally distant exons resulted in PCR amplification across the deleted region, allowing precise identification of the sequence breakpoint (Fig. 3 and 4 in the Supplementary Appendix). The deletion-spanning PCR products were not amplified from the 38 patients with other immune disorders, including the 6 patients with autosomal dominant HIES, or from the 115 healthy control subjects.

Heterozygous deletions in the *DOCK8* gene



**Figure 2. Immunologic Assessment.**

Values for blood analyses are shown according to age in the index patients in Families 1, 2, and 3; Patients 1 and 2 in Family 4; Patients 1 and 2 in Family 5; Patient 1 in Family 6; Patient 1 in Family 7; and Patients 1 and 2 in Family 8. In the color key, patients are listed according to their family number, followed by their patient number (Patient 1-1, Patient 2-1, etc.). Normal ranges for age are shown in gray. For lymphocyte subgroups, the normal ranges represent the 10th to 90th percentiles based on published data.<sup>20</sup> For leukocyte subgroups and serum immunoglobulin, the normal ranges are shown with 95% confidence intervals.<sup>21</sup> For eosinophils, the lines represent arbitrary thresholds for mild, moderate, or severe eosinophilia.<sup>22</sup>


were found in patients from Families 3, 4, 5, and 6: deletions C, D, E, and F, respectively (Fig. 2A and 2C in the Supplementary Appendix). As expected, the heterozygous deletions resulted in apparent homozygosity of single-nucleotide polymorphisms (SNPs) within the corresponding sequenced regions (data not shown). Deletions A through F were novel structural variants that were not found in the Database of Genomic

Variants, and no homozygous losses in this area of copy-number variation have been reported previously.

Patients with heterozygous deletions in one allele of *DOCK8* had compound heterozygosity for deleterious mutations (Fig. 2B and 2C in the Supplementary Appendix). None of these mutations were reported in SNP databases, and they were not identified in the 38 patients with other

**Figure 3. Patient Pedigrees and *DOCK8* Molecular Analyses.**

Panel A shows the pedigrees of the eight families in the study. Proband is designated by arrows. Circles represent female family members, and squares represent male family members. A slash through a symbol represents a deceased person. Panel B shows log<sub>2</sub> ratios of DNA from Patient 1 of Family 1 with deletion A (spanning exons 10 through 23) to reference DNA (on the x axis) after hybridization to probes at or near the *DOCK8* locus, according to the genomic position (on the y axis). The arrow pointing to the shaded region indicates the homozygous *DOCK8* deletion. The green crosses indicate probes where the signal of the log<sub>2</sub> ratio was less than zero, and the orange crosses more than zero. Panel C shows immunoblots for *DOCK8* protein expression in seven families that were included in the study. Lysates are from primary T cells (for all members of Family 1 and Patient 2 of Family 8), Epstein-Barr virus-transformed B-cell lines (for Families 3, 4, 6, and 7 and Patient 1 of Family 8), and herpesvirus saimiri-transformed T-cell lines (for Family 5). The data for Family 6 show two lanes (samples from a control [C] and from Patient 1) that are aligned from the same gel. Immunoblots for Patient 1 in Family 2 and for Patient 1 in Family 5, which were analyzed after the completion of the study, are shown in Figure 7A in the Supplementary Appendix. No *DOCK8* protein was found in samples from any of the 11 patients who were tested. Included in the samples for Family 1 is an unrelated *STAT3*-mutant control with autosomal dominant hyperimmunoglobulin E syndrome.  $\beta$ -actin is included as a loading control in all the analyses. P denotes parent, and S a heterozygous member of Family 4.



immune disorders, including the 6 patients with autosomal dominant HIES, or in the 115 control subjects.

Overall, the genetic variants consisted of large missing portions of the *DOCK8* coding sequence, including a conserved DOCK homology region 1 (DHR1) domain, or had other expected harmful effects in the gene (Fig. 2C in the Supplementary Appendix). *DOCK8* messenger RNA (mRNA) is present in tissues from lung, kidney, pancreas, and placenta, but it is unknown whether hematopoietic cells express *DOCK8*.<sup>16</sup> We found that monocytes, B cells, and T cells from healthy blood donors contain *DOCK8* mRNA, as assessed by quantitative reverse-transcriptase-PCR (qRT-PCR) assay; high levels were found in activated, expanded primary T-cell cultures and transformed lymphocyte lines (Fig. 6A in the Supplementary Appendix). Immunoblotting showed *DOCK8* protein in lymphocytes from an unrelated patient

with autosomal dominant HIES (Fig. 3C). By contrast, DOCK8 protein was not detected in primary T-cell cultures or transformed lymphocyte lines from all 11 patients who were tested (Fig. 3C, and Fig. 7A in the Supplementary Appendix). Residual levels of full-length protein in lymphocytes from Patient 2 in Family 8 were detected on overexposure of the immunoblot. Lymphocytes from Patient 1 in Family 1 contained minimal levels of a truncated protein that had been generated from the remaining exons (Fig. 7B in the Supplementary Appendix). These results are consistent with the finding of minimal amounts of transcripts or none in the patients' lymphocytes (Fig. 6B in the Supplementary Appendix). In the case of premature stop codons, this finding is most likely due to nonsense-mediated decay.<sup>23</sup>

#### FUNCTION OF CD8 T CELLS

CD8 T cells did not expand well from activated DOCK8-deficient peripheral-blood mononuclear cells from Patient 1 in Family 1 (Fig. 4A) or from three additional patients (Fig. 8A in the Supplementary Appendix). This result was not due to a low starting number of CD8 T cells, since the number of cells did not increase when they were normalized to the starting number. Moreover, CFSE dilution, a marker of cell division, was impaired in DOCK8-deficient CD8 T cells after stimulation of T-cell receptors with anti-CD3 plus anti-CD28 antibodies (Fig. 4B, and Fig. 8C in the Supplementary Appendix). By contrast, we observed no defect in the proliferation of CD4 T cells (Fig. 8B, 8C, and 8D in the Supplementary Appendix).


The induction of cell-surface activation marker CD25 on CD8 T cells was impaired in two of three patients who were tested (Fig. 9A in the Supplementary Appendix). The expression of the antiviral cytokines interferon- $\gamma$  and tumor necrosis factor  $\alpha$ , as assessed by intracellular flow cytometry, was moderately decreased in CD8 T cells in two patients who were tested (Fig. 9B in the Supplementary Appendix). Measurements of cytotoxicity in CD8 T cells on a per-cell basis in four patients were similar to those in control subjects. Both intracellular perforin and degranulation, as assessed by cell-surface extravasation of lysosomal-associated membrane protein 1 (LAMP1, also called CD107a), were normal (Fig. 9C and 9D in the Supplementary Appendix).

#### DISCUSSION

DOCK8 is a member of the DOCK180 superfamily of guanine nucleotide exchange factors, which interact with Rho GTPases.<sup>16</sup> The function of DOCK8 is unknown, but by analogy to related molecules, the protein probably regulates cytoskeletal rearrangements that are required for cellular structure, migration, adhesion, and other functions.<sup>24</sup> We found germ-line loss-of-function DOCK8 mutations in patients with a variant of combined immunodeficiency characterized by unusual susceptibility to cutaneous viral infections and cancers. In addition, DOCK8 deficiency was found in a subgroup of patients who were previously thought to have autosomal recessive HIES with severe allergic manifestations. Allergy is not characteristic of other forms of primary immunodeficiency diseases associated with elevated IgE levels, including autosomal dominant HIES.<sup>15</sup>

The immunologic phenotype in patients with DOCK8 deficiency resembles certain features of mice that have been rendered genetically deficient in a related gene, *dock2*, by homologous recombination. Such mice have a decreased number of T cells, reduced responsiveness of the T-cell antigen receptor, and allergic disease with high levels of serum IgE.<sup>25-27</sup> However, unlike patients with DOCK8 deficiency, *dock2*-deficient mice have no reported susceptibility to experimental viral infections. Although abnormal neutrophil chemotaxis was found in two of four patients who were tested (data not shown), it has not yet been determined whether the patients had the types of impaired lymphocyte chemotaxis and abnormal lymphoid organ architecture that have been seen in mice.<sup>25,28</sup> We evaluated the patients with immunodeficiency for a deficiency of the human orthologue of *dock2* but found no evidence of such a deficiency (data not shown). We cannot rule out cooperative effects, but our findings suggest that DOCK2 and DOCK8 do not serve redundant functions in humans. Moreover, the independent discovery of *dock8* mutations as a cause of immunodeficiency in ethylnitrosourea-mutagenized mice complements our findings (Cornall R, Goodnow C; personal communication).

HIES consists of primary immunodeficiencies that are characterized by severe eczema, recurrent skin infections (often from *S. aureus*), mucocuta-



**Figure 4. Impaired CD8 T-Cell Activation and Proliferation in Association with *DOCK8* Deficiency in Family 1.**

Panels A and B show the results of two independent experiments in Family 1. In Panel A, after activation of peripheral-blood mononuclear cells (PBMCs) with anti-CD3 plus anti-CD28 antibodies for 3 days, T cells were expanded in culture with interleukin-2. Absolute numbers of CD8 T cells were calculated on the basis of flow cytometric determinations of their percentages; CD8 T-cell numbers over time are shown as factor increases normalized to the prestimulated cell population. In Panel B, PBMCs that were labeled with carboxy-fluorescein succinimidyl ester (CFSE) were either unstimulated or stimulated, as described for Panel A. After 3 days, cells were stained with the use of fluorescence-labeled antibodies against T-cell subgroup markers and then analyzed on flow cytometry. The top row shows profiles for stimulated cells, with the location of each dot reflecting an individual cell's intensity for CD4 staining (on the x axis) versus CD8 staining (on the y axis). The boxes delineate the CD8 and CD4 T-cell subgroups. As compared with control samples, samples from Patient 1 had a lower proportion of CD8 T cells (12% of lymphocytes) after activation. Gated CD8 T cells were further analyzed for the fluorescence intensity of CFSE, which is successively diluted with each cell division. The bottom row shows histograms with the height (y axis) reflecting a percent of maximum of gated CD8 T cells that display any given fluorescence intensity of CFSE (on the x axis). The gray shading indicates unstimulated cells, and the blue line indicates stimulation. C denotes healthy control subject, and P parent.

neous candidiasis, recurrent sinopulmonary infections, elevated serum IgE levels, and eosinophilia.<sup>6</sup> Autosomal dominant HIES results from dominant interfering mutations in the gene encoding transcription factor STAT3. Such mutations lead to failed differentiation of a specialized subgroup of helper T cells that produce interleukin-17 for defense against fungal and extracellular bacterial infections.<sup>18,29-32</sup> By contrast, the autosomal recessive form features recurrent cutaneous viral infections but not pneumatocele formation or connective-tissue and skeletal abnormalities.<sup>5</sup> A homozygous loss-of-function mutation in the tyrosine kinase 2 gene, which en-

codes the TYK2 transcription factor, resulted in impaired cytokine signaling in multiple pathways in a single patient with elevated IgE levels but not in other patients.<sup>33,34</sup> Patients with these forms of HIES characteristically have a normal number of lymphocytes.<sup>5,6,33</sup>

The *DOCK8* immunodeficiency syndrome accounts for subgroups of patients who have undefined combined immunodeficiencies or who were previously misclassified as having autosomal recessive HIES. Besides lymphopenia, *DOCK8* deficiency has manifestations that have not previously been associated with HIES, including cancers related to cutaneous viral infections. Squamous-



cell cancers occurred in three of our patients who had extensive human papillomavirus infections, and one patient also had cutaneous T-cell lymphoma–leukemia. Impaired CD8 T cells in DOCK8 deficiency suggests that these cancers may be due to impaired tumor surveillance. It is also possible that DOCK8 has tumor-suppressor functions. Indeed, *DOCK8* deletions in primary lung cancers, gastric- and breast-cancer lines, and gliomas suggest a role of DOCK8 in tumor suppression.<sup>35,36</sup> Heterozygous *DOCK8* deletions have been reported in several cases of mental retardation, developmental delay, and autistic-spectrum disorder.<sup>37,38</sup> However, most of the DOCK8-deficient patients and their heterozygous relatives in our study had normal intelligence, suggesting that DOCK8 is not essential for cognitive function.

Other heterozygous deletions and duplications, reported in the Database of Genomic Variants, are consistent with the copy-number variation of DOCK8 within the normal population. Our findings support the increasing recognition that these and other structural variants, which have been estimated to account for a least 5% of the hu-

man genome and are enriched in genes of the immune system, represent sources of variation that can contribute to mendelian human diseases and influence immune function with respect to susceptibility to infections.<sup>39</sup>

Supported by the Intramural Research Program of the National Institutes of Health, the National Institute of Allergy and Infectious Diseases, and the National Cancer Institute, and by a grant from the Howard Hughes Medical Institute–National Institutes of Health Research Scholar Program (to Mr. Davis). Sequencing was performed by the Genomics Unit of the Rocky Mountain Laboratories Research Technologies Section of the National Institute of Allergy and Infectious Diseases.

No potential conflict of interest relevant to this article was reported.

We thank Drs. John O'Shea, Pamela Schwartzberg, Andrew Snow, Harry Malech, and Michael Lenardo for helpful discussions or critical reading of the manuscript; Dr. Steve Porcella and Kent Barbian of the Rocky Mountain Laboratories Genomics Unit for sequencing support; Dr. Jennifer Johnston and Amy Hsu-Lin for technical advice; Christopher Dove and Timothy Lenardo for experimental assistance; Drs. Aino Ruusala and Pontus Aspenström for reagents; Christine Spalding for clinical support; Drs. Prescott Atkinson, Michael Cooperstock, Carolyn Fein-Levy, Robin La-Croix, Joe LaRussa, Harold Lischner, and Swayam Sadanand for patient referrals; Drs. Jennifer Puck and Bodo Grimbacher for their past contributions in assembling several of the case reports of HIES; Andy Johnson and Drs. Michael Lenardo, Richard Cornell, and Christopher Goodnow for sharing information about their independent and complementary findings with respect to the consequences of *dock8* mutations in mice; and the patients and their families for participating in the study.

## REFERENCES

1. Fischer A. Human primary immunodeficiency diseases. *Immunity* 2007;27:835-45.
2. Bustamante J, Zhang SY, von Bernuth H, Abel L, Casanova JL. From infectious diseases to primary immunodeficiencies. *Immunol Allergy Clin North Am* 2008;28:235-58.
3. Kawai T, Malech HL. WHIM syndrome: congenital immune deficiency disease. *Curr Opin Hematol* 2009;16:20-6.
4. Prawer SE, Pass F, Vance JC, Greenberg LJ, Yunis EJ, Zelikson AS. Depressed immune function in epidermodysplasia verruciformis. *Arch Dermatol* 1977;113:495-9.
5. Renner ED, Puck JM, Holland SM, et al. Autosomal recessive hyperimmunoglobulin E syndrome: a distinct disease entity. *J Pediatr* 2004;144:93-9.
6. Freeman AF, Holland SM. The hyper-IgE syndromes. *Immunol Allergy Clin North Am* 2008;28:277-91.
7. Kalman L, Lindegren ML, Kobrynski L, et al. Mutations in genes required for T-cell development: IL7R, CD45, IL2RG, JAK3, RAG1, RAG2, ARTEMIS, and ADA and severe combined immunodeficiency: HuGE review. *Genet Med* 2004;6:16-26.
8. van der Burg M, van Veelen LR, Verkaik NS, et al. A new type of radiosensitive T-B-NK+ severe combined immunodeficiency caused by a LIG4 mutation. *J Clin Invest* 2006;116:137-45.
9. van der Burg M, Ijspeert H, Verkaik NS, et al. A DNA-PKcs mutation in a radiosensitive T-B- SCID patient inhibits Artemis activation and nonhomologous end-joining. *J Clin Invest* 2009;119:91-8.
10. Ahnesorg P, Smith P, Jackson SP. XLF interacts with the XRCC4-DNA ligase IV complex to promote DNA nonhomologous end-joining. *Cell* 2006;124:301-13.
11. Markert ML, Hershfield MS, Schiff RI, Buckley RH. Adenosine deaminase and purine nucleoside phosphorylase deficiencies: evaluation of therapeutic interventions in eight patients. *J Clin Immunol* 1987;7:389-99.
12. Pignata C, Fiore M, Guzzetta V, et al. Congenital alopecia and nail dystrophy associated with severe functional T-cell immunodeficiency in two sibs. *Am J Med Genet* 1996;65:167-70.
13. Shiow LR, Roadcap DW, Paris K, et al. The actin regulator coronin 1A is mutant in a thymic egress-deficient mouse strain and in a patient with severe combined immunodeficiency. *Nat Immunol* 2008;9:1307-15.
14. Liston A, Enders A, Siggs OM. Unravelling the association of partial T-cell immunodeficiency and immune dysregulation. *Nat Rev Immunol* 2008;8:545-58.
15. Ozcan E, Notarangelo LD, Geha RS. Primary immune deficiencies with aberrant IgE production. *J Allergy Clin Immunol* 2008;122:1054-62.
16. Ruusala A, Aspenström P. Isolation and characterisation of DOCK8, a member of the DOCK180-related regulators of cell morphology. *FEBS Lett* 2004;572:159-66.
17. Lei JY, Wang Y, Jaffe ES, et al. Microcystic adnexal carcinoma associated with primary immunodeficiency, recurrent diffuse herpes simplex virus infection, and cutaneous T-cell lymphoma. *Am J Dermatopathol* 2000;22:524-9.
18. Milner JD, Brenchley JM, Laurence A, et al. Impaired T(H)17 cell differentiation in subjects with autosomal dominant hyper-IgE syndrome. *Nature* 2008;452:773-6.
19. Grimbacher B, Schäffer AA, Holland SM, et al. Genetic linkage of hyper-IgE syndrome to chromosome 4. *Am J Hum Genet* 1999;65:735-44.
20. Shearer WT, Rosenblatt HM, Gelman RS, et al. Lymphocyte subsets in healthy children from birth through 18 years of age: the Pediatric AIDS Clinical Trials Group P1009 study. *J Allergy Clin Immunol* 2003;112:973-80.
21. Custer JW, Rau RE, eds. The Harriet Lane handbook: a manual for pediatric house officers. 18th ed. Philadelphia: Elsevier Mosby, 2009.
22. Brito-Babapulle F. The eosinophilias, including the idiopathic hypereosinophilic syndrome. *Br J Haematol* 2003;121:203-23.
23. Chang YF, Imam JS, Wilkinson MF.

The nonsense-mediated decay RNA surveillance pathway. *Annu Rev Biochem* 2007;76:51-74.

24. Meller N, Merlot S, Guda C. CZH proteins: a new family of Rho-GEFs. *J Cell Sci* 2005;118:4937-46.

25. Fukui Y, Hashimoto O, Sanui T, et al. Haematopoietic cell-specific CDM family protein DOCK2 is essential for lymphocyte migration. *Nature* 2001;412:826-31.

26. Sanui T, Inayoshi A, Noda M, et al. DOCK2 is essential for antigen-induced translocation of TCR and lipid rafts, but not PKC-theta and LFA-1, in T cells. *Immunity* 2003;19:119-29.

27. Tanaka Y, Hamano S, Gotoh K, et al. T helper type 2 differentiation and intracellular trafficking of the interleukin 4 receptor-alpha subunit controlled by the Rac activator Dock2. *Nat Immunol* 2007;8:1067-75.

28. Nombela-Arrieta C, Mempel TR, Soriano SF, et al. A central role for DOCK2 during interstitial lymphocyte motility and sphingosine-1-phosphate-mediated egress. *J Exp Med* 2007;204:497-510.

29. Holland SM, DeLeo FR, Elloumi HZ, et al. STAT3 mutations in the hyper-IgE syndrome. *N Engl J Med* 2007;357:1608-19.

30. Minegishi Y, Saito M, Tsuchiya S, et al. Dominant-negative mutations in the DNA-binding domain of STAT3 cause hyper-IgE syndrome. *Nature* 2007;448:1058-62.

31. Renner ED, Rylaarsdam S, Anover-Sombke S, et al. Novel signal transducer and activator of transcription 3 (STAT3) mutations, reduced T(H)17 cell numbers, and variably defective STAT3 phosphorylation in hyper-IgE syndrome. *J Allergy Clin Immunol* 2008;122:181-7.

32. de Beaucoudrey L, Puel A, Filipe-Santos O, et al. Mutations in STAT3 and IL12RB1 impair the development of human IL-17-producing T cells. *J Exp Med* 2008;205:1543-50.

33. Minegishi Y, Saito M, Morio T, et al. Human tyrosine kinase 2 deficiency reveals its requisite roles in multiple cytokine signals involved in innate and acquired immunity. *Immunity* 2006;25:745-55.

34. Woellner C, Schäffer AA, Puck JM, et al. The hyper IgE syndrome and mutations in TYK2. *Immunity* 2007;26:535.

35. Takahashi K, Kohno T, Ajima R, et al. Homozygous deletion and reduced expression of the DOCK8 gene in human lung cancer. *Int J Oncol* 2006;28:321-8.

36. Idhah A, Carvalho Silva R, Crinière E, et al. Genomic changes in progression of low-grade gliomas. *J Neurooncol* 2008;90:133-40.

37. Griggs BL, Ladd S, Saul RA, DuPont BR, Srivastava AK. Dedicator of cytokinesis 8 is disrupted in two patients with mental retardation and developmental disabilities. *Genomics* 2008;91:195-202.

38. Vinci G, Chantot-Bastaraud S, El Houate B, Lortat-Jacob S, Brauner R, McElreavey K. Association of deletion 9p, 46,XY gonadal dysgenesis and autistic spectrum disorder. *Mol Hum Reprod* 2007;13:685-9.

39. Sharp AJ, Cheng Z, Eichler EE. Structural variation of the human genome. *Annu Rev Genomics Hum Genet* 2006;7:407-42.

Copyright © 2009 Massachusetts Medical Society.

#### FULL TEXT OF ALL JOURNAL ARTICLES ON THE WORLD WIDE WEB

Access to the complete contents of the *Journal* on the Internet is free to all subscribers. To use this Web site, subscribers should go to the *Journal's* home page ([NEJM.org](http://NEJM.org)) and register by entering their names and subscriber numbers as they appear on their mailing labels. After this one-time registration, subscribers can use their passwords to log on for electronic access to the entire *Journal* from any computer that is connected to the Internet. Features include a library of all issues since January 1993 and abstracts since January 1975, a full-text search capacity, and a personal archive for saving articles and search results of interest. All articles can be printed in a format that is virtually identical to that of the typeset pages. Beginning 6 months after publication, the full text of all Original Articles and Special Articles is available free to nonsubscribers.